

Nuclear uncertainties in the NeNa-MgAl cycles and synthesis of ^{22}Na and ^{26}Al in classical novae

JORDI JOSÉ^{1,2}, ALAIN COC³, MARGARITA HERNANZ²

¹ *Departament de Física i Enginyeria Nuclear (UPC), Av. Víctor Balaguer s/n, E-08800 Vilanova i la Geltrú (Barcelona), SPAIN*

² *Institut d'Estudis Espacials de Catalunya (IEEC/CSIC/UPC), Edifici Nexus-201, Gran Capità 2-4, E-08034 Barcelona, SPAIN*

³ *Centre de Spectrométrie Nucléaire et de Spectrométrie de Masse, IN2P3-CNRS, Bât.104, F-91140 Orsay Campus, FRANCE*

Abstract. Classical novae eject significant amounts of matter into the interstellar medium, as a result of thermonuclear runaways. Nucleosynthesis associated with nova outbursts includes products from explosive H-burning, such as ^{17}O , ^{15}N and ^{13}C , and also radioactive species like ^7Be , ^{22}Na and ^{26}Al . In this paper we report on new hydrodynamic calculations of nova outbursts, from the onset of accretion to mass ejection. We stress the role played by the nuclear uncertainties associated with key reactions of the NeNa-MgAl cycles on the synthesis of ^{22}Na and ^{26}Al .

1 Introduction

The thermonuclear runaway model has been successful in reproducing the main observational properties of classical nova outbursts. According to this scenario, novae are produced by thermonuclear runaways located in the white dwarf component of a close binary system. The large, main sequence companion overfills its Roche lobe, providing matter outflows through the inner Lagrangian point that lead to the formation of an accretion disk around the white dwarf. A fraction of this H-rich matter lost by the companion ultimately ends up on top of the white dwarf, where it is gradually compressed as accretion goes on. The piling up of matter heats the envelope up to the point when ignition conditions to drive a TNR are reached.

An extended set of hydrodynamic computations of classical nova outbursts has been performed during the last 25 years (see [25], [17], [15], and references therein), for a wide range of white dwarf masses and initial luminosities, which have been able to identify two types of nova outbursts, those occurring in CO and in ONe white dwarfs. The latter ones have provided a framework for the origin of the high concentrations of Ne and more massive isotopes found in the spectra of some well-observed nova systems [19]. Among the isotopes synthesized in these so-called ONe novae (mainly ^{13}C , ^{17}O and ^{15}N), two radioactive species have raised a particular astrophysical interest: ^{22}Na and ^{26}Al . In this paper we investigate the role played by the nuclear uncertainties associated with key reactions of the NeNa-MgAl cycles on the synthesis of ^{22}Na and ^{26}Al during classical nova outbursts.

2 Synthesis of ^{22}Na and ^{26}Al in nova outbursts

The role of ^{22}Na as a diagnosis of a nova outburst was first suggested by [4]. In its decay to a short-lived excited state of ^{22}Ne ($\tau \sim 3.75$ yr), ^{22}Na emits a γ -ray photon of 1.275 MeV. Through this mechanism, nearby ONe novae within a few kiloparsecs

from the Sun may provide detectable γ -ray fluxes. Several experimental verifications of this γ -ray emission at 1.275 MeV from nearby novae have been attempted in the last twenty years, from which upper limits on the ejected ^{22}Na have been derived. In particular, the observations performed with the COMPTEL experiment on-board CGRO of five recent neon-type novae [12], as well as observations of *standard* novae, have led to an upper limit of $3.7 \times 10^{-8} M_{\odot}$ for the ^{22}Na mass ejected by any nova in the Galactic disk. This limit has posed some constraints on pre-existing theoretical models of classical nova explosions.

^{26}Al is another unstable nucleus, with a lifetime of $\tau = 1.04 \times 10^6$ years, that decays from ground state to the first excited level of ^{26}Mg , which in turn de-excites to its ground state by emitting a gamma-ray photon of 1.809 MeV. This characteristic gamma-ray signature, first detected in the Galactic Center by the HEAO-3 satellite, has been confirmed by other space missions. The most recent measurements made with COMPTEL have provided a map of the 1.809 MeV emission in the Galaxy ([6], [22]). The inferred $1 - 3 M_{\odot}$ of Galactic ^{26}Al are, according to the observed distribution, mainly attributed to young progenitors, such as massive AGB stars, SN II and Wolf-Rayet stars. However, a potential contribution from novae or low-mass AGB stars cannot be ruled out.

First estimates of the ^{22}Na and ^{26}Al production in novae were performed by different groups, using simplified one-zone models with representative temperature and density profiles (see [10], [31], [30], [20]), on the basis of ONeMg white dwarf stars. This scenario was revisited by [21] using hydrodynamic computations: the obtained ^{22}Na yields range between 5×10^{-5} and 5×10^{-3} , by mass. Assuming that the whole envelope ($\sim 10^{-4} - 10^{-5} M_{\odot}$) is ejected during the outburst, they derive peak fluxes of the ^{22}Na 1.275 MeV line in the range $(2 - 30) \times 10^{-5}$ photons $\text{s}^{-1} \text{cm}^{-2}$ (corresponding to ONeMg white dwarf masses between $1.00 - 1.35 M_{\odot}$, respectively) for classical novae exploding at 1 kpc. Their main conclusion, based on the reported 3σ line sensitivity of OSSE and COMPTEL, is that nearby classical novae involving ONeMg white dwarfs of $M_{wd} \geq 1.25 M_{\odot}$ should produce detectable ^{22}Na γ -rays for CGRO, a prediction not confirmed so far [12]. Their results showed also a significant production of ^{26}Al (i.e., $(19.6 - 7.5) \times 10^{-3}$, by mass, for the same white dwarf mass range), which could account for a major fraction of the Galactic ^{26}Al .

Recent hydrodynamic computations ([16], [14], [15]) using updated nuclear reaction rates, have led to a significant reduction of both ^{26}Al and ^{22}Na ejected during nova outbursts. In particular, a mean mass fraction of 1×10^{-4} of ^{22}Na is found in the $1.25 M_{\odot}$ ONe Model (with $M_{ejec}(^{22}\text{Na}) = 1.3 \times 10^{-9} M_{\odot}$), whereas a maximum value of 6×10^{-4} results from the $1.35 M_{\odot}$ ONe Model (with $M_{ejec}(^{22}\text{Na}) = 2.6 \times 10^{-9} M_{\odot}$). The corresponding peak fluxes in the 1.275 MeV ^{22}Na line, below 10^{-5} counts $\text{s}^{-1} \text{cm}^{-2}$ for novae at 1 kpc, turn out to be too low to be detected with OSSE or COMPTEL but represent potential targets for the nearby future INTEGRAL mission [7]. Concerning ^{26}Al , yields ranging from 2×10^{-3} to 2×10^{-4} by mass have been obtained in a series of ONe nova models with masses between $1.15 - 1.35 M_{\odot}$. Contribution of novae to the Galactic ^{26}Al is limited to $0.1 - 0.4 M_{\odot}$ [16].

Other hydrodynamic computations performed by [24] and [25], using also updated nuclear reaction rates and opacities, have modified previous estimates (i.e., [21]). The expected abundance of ^{22}Na in the ejecta has risen up to $(2 - 3) \times 10^{-3}$, by mass, when $1.25 M_{\odot}$ ONeMg white dwarfs are adopted, high enough to be detected by

CGRO, provided that all the accreted envelope ($\sim 3 \times 10^{-5} M_{\odot}$) is ejected. On the other hand, the improved input physics translates into a factor of 10 reduction on the synthesis of ^{26}Al , in better agreement with the analysis of the 1.809 MeV emission map provided by COMPTEL, and also with the results previously reported by [16].

Since the synthesis of both ^{22}Na and ^{26}Al is very dependent on the adopted nuclear reaction rates, we have investigated the effect of the nuclear uncertainties accompanying reaction rates within the NeNa-MgAl cycles on the synthesis of ^{22}Na and ^{26}Al , using an updated version of the spherically symmetric, hydrodynamic SHIVA code [15]. In a first step, we have identified the most relevant reactions involved in the synthesis of such isotopes. Main results are summarized in Table 1. We stress the crucial role played by $^{21}\text{Na}(p,\gamma)^{22}\text{Mg}$: a reduction by a factor 100 of the rate given by [1] translates into an increase in the mean abundance of ^{22}Na in the ejecta by a factor between 1.2 and 3. An increase by a factor of 3 is also obtained when the rate given by [26], instead of that from [1], is adopted for $^{22}\text{Na}(p,\gamma)^{23}\text{Mg}$. Worth noticing is also the increase in the final amount of ^{26}Al by a factor 2 when the term corresponding to the 0.188 MeV resonance, as measured by [29], is reduced by just a factor 1/3.

Table 1. Test Models

Nuclear reaction	Old rate	Test rate	$\frac{X(^{22}\text{Na})_{\text{test}}}{X(^{22}\text{Na})_{\text{old}}}$	$\frac{X(^{26}\text{Al})_{\text{test}}}{X(^{26}\text{Al})_{\text{old}}}$
1.15 $M_{\odot}$ ONe				
$^{21}\text{Na}(p,\gamma)^{22}\text{Mg}$	CF88	CF88/100	2.3	1.1
$^{25}\text{Al}(p,\gamma)^{26}\text{Si}$	Wie86	Coc95, case A	1	1.2
	Wie86	Coc95, case C	1	0.5
$^{23}\text{Na}(p,\gamma)^{24}\text{Mg}$	CF88	CF88+Gor89	1.2	1.3
$^{26}\text{Al}^g(p,\gamma)^{27}\text{Si}$	Vog89	Vog89	1	1.9
1/3×res(0.188 MeV)				
$^{26}\text{Al}^m(p,\gamma)^{27}\text{Si}$	CF88	CF88×100	1	1
$^{26}\text{Si}(p,\gamma)^{27}\text{P}$	Wie86	Her95	1	1
$^{27}\text{Al}(p,\alpha)^{24}\text{Mg}$	Cha88	Cha88+Tim88	1	1
$^{26}\text{Mg}(p,\gamma)^{27}\text{Al}$	Ili90	Ili90+Cha90	1	1
1.25 $M_{\odot}$ ONe				
$^{21}\text{Na}(p,\gamma)^{22}\text{Mg}$	CF88	CF88/100	3	1.1
$^{22}\text{Na}(p,\gamma)^{23}\text{Mg}$	CF88	Ste96	3	1.2
$^{23}\text{Mg}(p,\gamma)^{24}\text{Al}$	Wie86	Kub95	1.1	1.1
1.35 $M_{\odot}$ ONe				
$^{21}\text{Na}(p,\gamma)^{22}\text{Mg}$	CF88	CF88/100	1.2	1
$^{23}\text{Al}(p,\gamma)^{24}\text{Si}$	Wor94	Sch97	1	1

In a second step, a series of hydrodynamic nova models has been computed assuming upper, recommended and lower estimates of the reaction rates, from which limits on the production of both ^{22}Na and ^{26}Al are derived (see [13] for details). Main results are summarized in Table 2. Large differences in the ^{22}Na and ^{26}Al yields have been obtained: in particular, a factor ~ 4 in the ^{22}Na production for the 1.15 $M_{\odot}$ ONe Model, and a factor ~ 7 in the case of ^{26}Al production (1.35 $M_{\odot}$ ONe Model) when either the upper or the lower estimates are adopted. These differences are too large to accurately predict the contribution of classical novae to the Galactic ^{26}Al , or to

determine the maximum distance to a classical novae for a potential detection of the 1.275 MeV ^{22}Na γ -ray line.

Table 2. Ranges of ^{22}Na and ^{26}Al production

Model	Network	$X(^{22}\text{Na})$	$X(^{26}\text{Al})$	$\frac{X(^{22}\text{Na})_{\text{upper}}}{X(^{22}\text{Na})_{\text{lower}}}$	$\frac{X(^{26}\text{Al})_{\text{upper}}}{X(^{26}\text{Al})_{\text{lower}}}$
1.15 $M_{\odot}$ ONe	Lower	1.2E-4	3.3E-4	3.8	3.6
	Upper	4.6E-4	1.2E-3		
1.25 $M_{\odot}$ ONe	Lower	1.9E-4	2.0E-4	3.2	4.4
	Upper	6.0E-4	8.7E-3		
1.35 $M_{\odot}$ ONe	Lower	7.7E-4	1.5E-4	1.8	7.3
	Upper	1.4E-3	1.1E-3		

Therefore, we stress the need of new nuclear physics experiments to reduce the uncertainties associated with some key reactions of the NeNa-MgAl cycles, in particular $^{21}\text{Na}(p,\gamma)^{22}\text{Mg}$, $^{22}\text{Na}(p,\gamma)^{23}\text{Mg}$ and $^{25}\text{Al}(p,\gamma)^{26}\text{Si}$. A verification of the yet unpublished values corresponding to the 0.188 MeV resonance of $^{26}\text{Al}^g(p,\gamma)^{27}\text{Si}$, measured by [29], is also recommended.

Acknowledgements. This research has been partially supported by the DGICYT (PB97-0983-C03-02; PB97-0983-C03-03), by the CICYT-P.N.I.E., by the PICS 319, and by a CIRIT grant.

References

- [1] Caughlan G.R., Fowler W.A., 1988 (CF88), ADNDT 40, 283
- [2] Champagne A.E., et al., 1988 (Cha88), Nuc. Phys. A487, 433
- [3] Champagne A.E., Magnus P.V., Smith M.S., Howard A.J., 1990 (Cha90), Nuc. Phys. A512, 317
- [4] Clayton D.D., Hoyle F., 1974, ApJ 187, L101
- [5] Coc A., et al., 1995 (Coc95), A&A 299, 479
- [6] Diehl R., et al., 1995, A&A 298, 445
- [7] Gómez-Gomar J., Hernanz M., José J., Isern J., 1998, MNRAS 296, 913
- [8] Görres J., Wiescher M., Rolfs C., 1989 (Gor89), ApJ 343, 365
- [9] Herndl H., et al., 1995 (Her95), Phys. Rev. C52, 1078
- [10] Hillebrandt W., Thielemann F.-K., 1982, ApJ 225, 617
- [11] Iliadis Ch., et al., 1990 (Ili90), Nuc. Phys. A512, 509
- [12] Iyudin A.F., et al., 1995, A&A 300, 422
- [13] José J., Coc A., Hernanz M., 1998, in preparation
- [14] José J., Hernanz M., 1997, Nuc. Phys. A621, 491
- [15] José J., Hernanz M., 1998, ApJ 494, 680
- [16] José J., Hernanz M., Coc A., 1997, ApJ 479, L55
- [17] Kovetz A., Prialnik D., 1997, ApJ 477, 356
- [18] Kubono S., Kajino T., Kato S., 1995 (Kub95), Nuc. Phys. A588, 521
- [19] Livio M., Truran J.W., 1994, ApJ 425, 797
- [20] Nofar I., Shaviv G., Starrfield S., 1991, ApJ 369, 440
- [21] Politano M., et al., 1995, ApJ 448, 807
- [22] Prantzos N., Diehl R., 1996, Phys. Rep. 267, 1
- [23] Schatz H., et al., 1997 (Sch97), Phys. Rev. Let. 79, 3845
- [24] Starrfield S., Truran J.W., Wiescher M.C., Sparks W.M., 1997, Nuc. Phys. A621, 495
- [25] Starrfield S., Truran J.W., Wiescher M.C., Sparks W.M., 1998, MNRAS 296, 502
- [26] Stegmüller F., et al., 1996 (Ste96), Nuc. Phys. A601, 168
- [27] Timmermann R., et al., 1988 (Tim88), Nuc. Phys. A477, 105
- [28] Van Wormer L., et al., 1994 (Wor94), ApJ 432, 326
- [29] Vogelaar R.B., 1989 (Vog89), Ph.D. Thesis, California Institute of Technology
- [30] Weiss A., Truran J.W., 1990, A&A 238, 178
- [31] Wiescher M., Görres J., Thielemann F.-K., Ritter H., 1986 (Wie86), A&A 160, 56